

NPS55-79-028

NAVAL POSTGRADUATE SCHOOL

Monterey, California



CHANNELS THAT COOPERATIVELY SERVICE A DATA
STREAM AND VOICE MESSAGES, II:
DIFFUSION APPROXIMATIONS

by

J. P. Lehoczky

and

D. P. Gaver

November 1979

Approved for public release; distribution unlimited.

Prepared for:

Naval Research
., VA 22217

FEDDOCS
D 208.14/2:NPS-55-79-028

Naval Postgraduate School
Monterey, California

Rear Admiral T. F. Dedman
Superintendent

Jack R. Borsting
Provost

This report was prepared by:

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-79-028	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Channels that Cooperatively Service a Data Stream and Voice Messages, II: Diffusion Approximations		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) J. P. Lehoczky and D. P. Gaver		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS 61152N;R000-01-10 N0001480WR0054
11. CONTROLLING OFFICE NAME AND ADDRESS Chief of Naval Research Arlington, VA 22217		12. REPORT DATE November 1979
		13. NUMBER OF PAGES 43
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Communication Data-Voice Communications Channels Diffusion Approximation Queues Probability Models		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A system of channels mutually accommodate both data and voice messages, voice having preemptive priority but being a loss system, and data being allowed to queue. Approximations to the data queue properties are derived.		

CHANNELS THAT COOPERATIVELY SERVICE
A DATA STREAM AND VOICE MESSAGES, II:
DIFFUSION APPROXIMATIONS*

J. P. Lehoczký
Carnegie-Mellon University
Pittsburgh, PA

D. P. Gaver
Naval Postgraduate School
Monterey, CA

1. INTRODUCTION

In this paper, the study of the behavior of an element of a communication system carrying both data and voice traffic is continued. In a previous paper, Gaver and Lehoczký (1979), a fluid flow approximation was developed to predict the characteristics of the data queue length for such a system. The fluid flow approach applied to the case in which the data service rate, η , was large compared with the voice service rate, μ ; for example, when $\eta/\mu \sim 10^4$. In this paper, a different approach is taken: a Wiener process approximation is developed. The accuracy of this approximation depends on the "heavy traffic" assumption, that is that the overall traffic intensity should approach unity from below. One need not make any assumption concerning the individual parameters. It follows that the diffusion approximation complements the fluid flow approximation to give a more complete picture of the behavior of voice-data communication systems.

*Research in part sponsored by ONR at Naval Postgraduate School N001480WR00067, and in part by NSF at Carnegie-Mellon University, ENG79 05526.

The communication system to be studied is essentially a part of the SENET network as described by Coviello and Vena (1975) or Barbacci and Oakley (1976). This network employs time-slotted frames. A certain portion of each frame is allocated to voice traffic, while any data traffic can use all remaining capacity including any left unused by voice. The voice traffic cannot use any unused data capacity and operates as a loss system.

We introduce probabilistic assumptions, conventional in many queueing studies. Voice traffic arrives according to a Poisson(λ) process, and each voice customer has an independent exponential(μ) service time. Data arrivals are governed by an independent Poisson(δ) process and exhibit independent exponential(η) service times. A total of c channels are reserved for exclusive use of data, while v channels can be used by both data and voice; however, voice pre-empts data. The assumptions imply that voice can be modelled as an M/M/v/v loss system, and the well-known "Erlang B" loss formula will give the loss rate. We focus our attention on the behavior of the data queue and seek to develop expressions for the steady-state distribution and the mean queue length. Many authors have studied such a system, including Halfin and Segal (1972), Halfin (1972), Fischer and Harris (1976), Bhat and Fischer (1976), Fischer (1977), Chang (1977), and Gaver and Lehoczky (1979). Only the latter paper gives approximations valid for the extreme but realistic case in which η/μ is large, and none give a Wiener approximation.

Rather than focus on this simple version of the problem, we wish to provide a somewhat more general analysis. The generalizations arise when one considers more complex voice service. Two types of generalizations are rated here.

In general, voice traffic is unlike data traffic. A voice communication is actually a series of bursts or talkspurts separated by silence. It is reasonable to model a conversation as a two-state Markov chain of alternating talk and silence. Exponential holding times are appropriate for talkspurt and silence lengths, see Brady (1965). During the periods of silence, the voice channel is available for data traffic. This extra channel capacity can serve to allow for increased data utilization, or to reduce the data queue length, or both. The voice process can be described by a Markov chain for which each state provides a description of the number of voice customers requesting service and the number actually using the channel. For each voice state one can compute the number of channels available for data, and thus the data transition rates.

A second generalization allows for the addition of extra voice channels by reducing the quality. One strategy available to provide adequate voice and data capacity is to assign voice users to certain acceptable quality voice channels, say 6 or 8 KBPS. When a specified number of voice users are in the system, new voice users are assigned to less acceptable channels, say with 2 or 4 KBPS. In this way, one has increased the number of voice users that can be serviced at the expense of the quality of the voice

transmissions. This situation can also be incorporated into our structure. One again defines voice states: each state provides a description of the voice channels in use. One can specify a generator Q for the Markov process over the states and use it to compute the channel availability for data for each state. To include this and other possible generalizations, we assume that the voice process is a continuous time Markov chain with finite state space and that it subordinates the data queue length process. We will work out formulas for the special case in which the voice is an M/M/v/v queueing system.

2. Diffusion Approximation Approach

We assume that the voice process $\{V(t), t \geq 0\}$ can be described by a Markov chain with state space $\{1, 2, \dots, N\}$. We let this Markov chain have generator $Q = (q_{ij})$, an $N \times N$ matrix, and stationary distribution π . An important special case occurs when voice operates as an M/M/v/v loss system in which case

$$\tilde{Q} = \mu \tilde{S} = \mu \begin{pmatrix} -\rho_V & \rho_V & & & & & & & & \\ & 1 & -(1+\rho_V) & \rho_V & & & & & & \\ & & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \\ & & & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \\ & & & & v-1 & -(v-1+\rho_V) & \rho_V & & & \\ & & & & & & v & -v & & \end{pmatrix} \quad (2.1)$$

and then $\pi = (\pi_0, \dots, \pi_v)$ with

$$\pi_i = \frac{\rho_V^i / i!}{\sum_{j=0}^v \rho_V^j / j!}, \quad \rho_V = \frac{\lambda}{\mu}.$$

Each voice state i gives a number of channels available for data, say c_i . From c_i one can compute the rate, r_i , at which the data queue increases or decreases. Here $r_i = \delta - \eta c_i$. We let

$$\tilde{R} = \begin{pmatrix} r_1 & & & 0 \\ & \ddots & & \\ & & c & \\ & & & r_N \end{pmatrix} \quad (2.2)$$

For stability one requires $\sum_{i=1}^N \pi_i r_i < 0$. In the special case for which \tilde{Q} is given by (2.1), these conditions become

$$\sum_{i=0}^v \frac{\rho_v^i}{i!} (\delta - (c + v - i)\eta) < 0$$

where data has exclusive use of c channels and the voice has v channels. This condition can be written $\rho < 1$ where $\rho = (\rho_d + \rho_v(1-q))/(c+v)$, $\rho_d = \delta/\eta$, $\rho_v = \lambda/\mu$, and q is the blocking probability

$$q = \frac{\rho_v^v/v!}{\sum_{j=0}^v \rho_v^j/j!}$$

We develop a Wiener process approximation under a heavy-traffic assumption that $\sum_{i=1}^N r_i \pi_i \nearrow 0$. This entails finding an appropriate infinitesimal mean and variance. Locally the data

queue length process does not resemble a Wiener process. If voice is in state i , the change in the data queue is nearly deterministic and equal to $r_i dt$ rather than a normally distributed random variable. Nevertheless, if we consider a longer period of time, say $[t, t+T]$ with T large, then the change in queue length will be more reasonably assumed to be normally distributed. The next section contains the outline of a proof that a suitably scaled version of the process converges to a Wiener process. This proof does not require the fluid flow approach of η/μ large, so a slightly more refined variance term is derived. The Wiener process derived will have a reflecting barrier at 0.

The voice process subordinates the data process; see Feller [], pp. for the notion of subordination. If voice is in state $V(t)$, then the data queue is increasing at rate $r_{V(t)}$ over $[0, t + dt]$. The expected change over $[t, t+T]$ is given by

$$E \int_t^{t+T} r_{V(s)} ds = \int_t^{t+T} E(r_{V(s)}) ds = \int_t^{t+T} \sum_{i=1}^N r_i P(v(s)=i) ds .$$

For the case of T large we can assume $V(t)$ has the stationary distribution, thus $V(s)$ has distribution π for all s . The expected change over $[t, t+T]$ thus becomes $T \sum_{i=1}^N r_i \pi_i$. Notice that the data queue is very large, so the boundary is not encountered. For a Wiener process, this expected change would be

mT again assuming that the boundary is not encountered. We thus find

$$m = \sum_{i=1}^N r_i \pi_i = \underline{\pi}^T \underline{R} \underline{1} \quad (2.3)$$

We follow a similar method to determine σ^2 . The variance in the change in the data queue length over $[t, t+T]$ is given by

$$\text{Var} \int_t^{t+T} r_{V(s)} ds .$$

We again assume $V(s)$ has distribution $\underline{\pi}$, so this variance is equal to

$$\begin{aligned} \text{Var} \int_0^T r_{V(s)} ds \\ = E \left(\int_0^T r_{V(s)} ds \right)^2 - \left(E \int_0^T r_{V(s)} ds \right)^2 = E \left(\int_0^T r_{V(s)} \right)^2 - (\underline{\pi} \underline{R} \underline{1})^2 T^2 \end{aligned}$$

The last term can be computed to be

$$\begin{aligned} E \left(\int_0^T r_{V(s)} ds \right)^2 &= E \int_0^T \int_0^T r_{V(s)} r_{V(t)} ds dt = \int_0^T \int_0^T E(r_{V(s)} r_{V(t)}) ds dt \\ &= 2 \int_0^T \int_0^T \sum_{i=1}^N r_i P(V(s)=i) E(r_{V(t)} | V(s)=i) ds dt \\ &= 2 \int_0^T \int_0^T \sum_{i=1}^N r_i \pi_i \sum_{j=1}^N r_j P(v(t)=j | V(s)=i) ds dt . \end{aligned}$$

We let $p_{ij}(u) = P(V(u)=j|V(0)=i)$ and $\underline{p}(u) = (p_{ij}(u))$.

The Kolmogorov forward equations give $\underline{p}(u) = \exp(\underline{Q}u)$ where $\exp(\underline{M}) = \underline{I} + \underline{M} + \underline{M}^2/2! + \dots$ for a square matrix \underline{M} . One can reexpress the second moment in matrix form as

$$\begin{aligned} E\left(\int_0^T r_{V(s)} ds\right)^2 &= 2 \int_0^T \int_0^t \underline{\pi} \underline{R} \exp(\underline{Q}(t-s)) \underline{R} \underline{1} ds dt \\ &= 2 \underline{\pi} \underline{R} \int_0^T \int_0^t \exp(\underline{Q}u) du dt \underline{R} \underline{1} . \end{aligned}$$

In view of the fact that \underline{Q} is singular, one can most easily carry out the required integration by introducing the eigenvalue decomposition $\underline{Q} = \underline{\Phi} \underline{D} \underline{\Psi}$ with

$$\underline{D} = \begin{pmatrix} \theta_1 & & & \\ & \theta_2 & & \\ & & \ddots & \\ & & & \theta_N \end{pmatrix}$$

a diagonal matrix of eigenvalues,

$$\underline{\Phi} = (\underline{\phi}_1, \underline{\phi}_2, \dots, \underline{\phi}_N), \quad \underline{\Psi} = (\underline{\pi}, \underline{\psi}_2, \dots, \underline{\psi}_N)^T,$$

the associated right- and left-eigenvectors. Here $\underline{\Phi} \underline{\Psi} = \underline{\Psi} \underline{\Phi} = \underline{I}$. Since \underline{Q} is irreducible and finite θ_i has a negative real part for $2 \leq i \leq N$. It follows that

$$\int_0^T \int_0^t \exp(\underline{Q}_u) du dt = \int_0^T \int_0^t \exp(\underline{D}u) du dt$$

$$\underline{\Psi} = 2\Phi \begin{pmatrix} T^2/2 & & & \\ & \alpha_2 & & \\ & & \ddots & \\ & & & \alpha_N \end{pmatrix} \underline{\Psi}$$

with $\alpha_i = \theta^{-i}(\theta^{-i}(e^{\theta_i T} - 1))$. Since θ_i has a negative real part $\alpha_i \sim -T/\theta_i$ as $T \rightarrow \infty$. For large T , the second moment can be written approximately as

$$2\pi^T \underline{R}\Phi \begin{pmatrix} T^2/2 & & & \\ & -T\theta_1^{-1} & & \\ & & \ddots & \\ & & & -T\theta_N^{-1} \end{pmatrix} \underline{\Psi R1}$$

thus the variance is given by

$$T^2 \pi^+ \underline{R} \begin{pmatrix} 1 & & & \\ & 0 & & \\ & & \ddots & \\ & & & 0 \end{pmatrix} \underline{\Psi R1} - 2T\pi^T \underline{R}\Phi \begin{pmatrix} 0 & & & \\ & \theta_2^{-1} & & \\ & & \ddots & \\ & & & \theta_N^{-1} \end{pmatrix} \underline{\Psi R1} - T^2 (\pi^+ \underline{R1})^2.$$

The first and last terms cancel to give

$$\text{Var}\left(\int_0^T r_{V(s)} ds\right) = -2\tilde{\pi}^T \tilde{R} \tilde{\Phi} \begin{pmatrix} 0 & & & \\ & \theta_2^{-1} & & \\ & & \ddots & \\ & & & \theta_N^{-1} \end{pmatrix} \tilde{\Psi} R 1 \quad (2.4)$$

For a Wiener process, this variance would be given by $T\sigma^2$.

We let

$$\tilde{Q}^- = \tilde{\Phi} \begin{pmatrix} 0 & & & \\ & \theta_2^{-1} & & \\ & & \ddots & \\ & & & \theta_N^{-1} \end{pmatrix} \tilde{\Psi} \quad (2.5)$$

so \tilde{Q}^- is a reflexive generalized inverse of \tilde{Q} obtained by reciprocating nonzero eigenvalues. We note that both $\tilde{Q}\tilde{Q}^-\tilde{Q} = \tilde{Q}$ and $\tilde{Q}^-\tilde{Q}\tilde{Q}^- = \tilde{Q}^-$ hold, thus we write

$$\frac{\sigma^2}{2} = -\tilde{\pi}^T \tilde{R} \tilde{Q}^- R 1 \quad (2.6)$$

$$m = \tilde{\pi}^+ R 1 < 0 .$$

The crucial parameter $2m/\sigma^2$ can thus be calculated.

We illustrate the formulas in (2.6) for the special case where \tilde{Q} is given by (2.1). For this case

$$\frac{\sigma^2}{2} = \frac{1}{\mu} \pi^T R \Phi \begin{pmatrix} 0 \\ \theta_1^{-1} \\ \cdot \\ \cdot \\ \cdot \\ \theta_v^{-1} \end{pmatrix} \Psi R 1 = \frac{1}{\mu} \pi^T R S^{-1} R 1$$

$$= \frac{1}{\mu} \pi^T R S^{-1} (R - mI) 1 + \frac{m}{\mu} \pi^T R S^{-1} 1 .$$

For our choice of S^{-1} , $S^{-1} 1 = 0$, thus

$$\frac{\sigma^2}{2} = \frac{1}{\mu} \pi^T R S^{-1} (R - mI) 1 = \frac{1}{\mu} \pi^T R S^{-1} y$$

where $y = (R - mI) 1$. We note that $\pi^T y = 0$, so if $z = S^{-1} y$, then $S z = y$ is a consistent set of equations and $\pi^T z = 0$.

In general one can solve (2.6) by direct calculation of Q^{-1} ; however, this is rarely possible. A second approach is to solve $Qz = y$ subject to $\pi^T z = 0$. For Q of form (2.1), this can be carried out. The equations become

$$\begin{aligned} \rho_v (z_i - z_{i-1}) - (i-1)(z_{i-1} - z_{i-2}) &= r_{i-1} - m, \quad 1 \leq i \leq v \\ -v(z_v - z_{v-1}) &= r_n - m \end{aligned} \quad (2.7)$$

$$\sum_{i=0}^v \frac{\rho_v^i}{i!} z_i = 0.$$

One can easily find

$$z_k - z_{k-1} = \frac{\sum_{j=0}^{k-1} \pi_j (r_j - m)}{\rho_v \pi_{k-1}} .$$

We define

$$T_k = \sum_{j=0}^k \pi_j (r_j - m) .$$

Thus $T_v = 0$ and $z_k - z_{k-1} = T_{k-1} / \rho_v \pi_{k-1}$. Moreover

$$z_k = z_0 + \sum_{j=1}^k z_j = z_0 + \sum_{j=1}^k \frac{T_{j-1}}{\rho_v \pi_{j-1}} .$$

The $\sum_{i=0}^v \pi_i z_i = 0$ condition gives

$$z_0 = - \sum_{k=1}^v \pi_k \sum_{j=1}^k \frac{T_{j-1}}{\rho_v \pi_{j-1}} .$$

The vector \tilde{z} has now been explicitly determined, and it remains to compute

$$\frac{1}{\mu} \tilde{\pi}^T \tilde{R} \tilde{z} = \frac{1}{\mu} \tilde{\pi}^T \tilde{R} (z_0 \mathbf{1} + (0, u_1, \dots, u_v))$$

with

$$u_k = \sum_{j=1}^k \frac{T_{j-1}}{\rho_v \pi_{j-1}} .$$

We find

$$\begin{aligned}
\frac{1}{\mu} \pi^T R z &= -\frac{m}{\mu} z_0 + \frac{1}{\mu} \sum_{k=1}^v \pi_k r_k \sum_{i=1}^k \frac{T_{i-1}}{\rho_v^{\pi_{i-1}}} \\
&= \frac{1}{\mu} \sum_{k=1}^v \pi_k (r_k^{-m}) \sum_{i=1}^k \frac{T_{i-1}}{\rho_v^{\pi_{i-1}}} \\
&= \frac{1}{\mu} \sum_{i=1}^v \frac{T_{i-1}}{\rho_v^{\pi_{i-1}}} \sum_{k=i}^v \pi_k (r_k^{-m}) = \frac{1}{\mu} \sum_{i=1}^v \frac{T_{i-1}}{\rho_v^{\pi_{i-1}}} \left(-\sum_{k=0}^{i-1} \pi_k (r_k^{-m}) \right) \\
&= -\frac{1}{\mu} \sum_{i=1}^v \frac{T_{i-1}^2}{\rho_v^{\pi_{i-1}}} = -\frac{1}{\mu} \sum_{i=0}^{v-1} \frac{T_i^2}{\rho_v^{\pi_i}} .
\end{aligned}$$

We thus derive

$$\frac{\sigma^2}{2} = \frac{1}{\mu} \sum_{i=0}^{v-1} \frac{T_i^2}{v_i} , \quad T_i = \sum_{j=0}^i \pi_j (r_j^{-m}) \quad (2.8)$$

For the special case considered $r_j = S - (c+v-j)\eta = \eta(\rho_d - (c+v) + j)$. One finds

$$\begin{aligned}
m &= \sum_{i=0}^v \eta(\rho_d - (c+v) + j) \frac{\rho_v^j}{j! S_v} \\
&= \eta[\rho_d - (c+v) + \rho_v(1-q)]
\end{aligned}$$

where $q = \rho_v^v / v! S_v$ is the blocking probability. Thus $m = -\eta(c+v)(1-\rho)$. Plugging into (2.8) we find

$$\frac{\sigma^2}{2} = \frac{\eta^2}{\rho_v \mu} \sum_{i=0}^{v-1} \frac{\left(\sum_{j=0}^i \frac{\rho_v^j}{j!} [\rho_d^{-(c+v)\rho} + j] \right)^2}{S_v \rho_v^i / i!} .$$

This can be simplified by letting $q_i = \rho_v^i / i! S_i$, the blocking probability for an M/M/i/i system with traffic intensity ρ_v .

We find

$$\frac{\sigma^2}{2} = \frac{\eta^2 \rho_v}{\mu} \sum_{i=0}^{v-1} \frac{S_i}{S_v} \frac{(q_i - q_v)^2}{q_i} , \quad S_i = \sum_{j=0}^i \frac{\rho_v^j}{j!} . \quad (2.9)$$

The special case $v = 1$ gives

$$\frac{\sigma^2}{2} = \frac{\eta^2}{\mu} \frac{\rho_v}{(1 + \rho_v)^2} .$$

We thus have given explicit formulas for the infinitesimal mean and variance of a Wiener process with reflecting barrier at 0. One can therefore use standard results for Wiener processes with reflecting barriers to derive the stationary distribution--it will be exponential with parameter $2m/\sigma^2$. The stationary mean queue length will thus be $\sigma^2/2m$; where σ^2 and m are given by (2.6) and (2.8). For example when $v = 1$ and $c = 0$

$$\frac{\sigma^2}{2m} = \frac{\eta}{\mu} \frac{\rho_v}{(1-\rho)(1+\rho_v)^3} .$$

This is developed under heavy traffic conditions, i.e., $\rho \nearrow 1$.

For such conditions

$$\rho_d + \frac{\rho_v}{1 + \rho_v} \rightarrow 1$$

or

$$\left| \rho_d - \frac{1}{1 + \rho_v} \right| \rightarrow 0 .$$

It follows that the above expression is also

$$\frac{\eta}{\mu} \frac{\rho_v \rho_d}{(1 - \rho)(1 + \rho_v)^2}$$

in agreement with Fischer (1977) and Gaver and Lehoczky (1979).

The characterization of the data queue as a Wiener process with reflecting barrier allows one to discuss the dynamics of the data queue. For example, suppose the data queue is at level x at time t where x is large. One might wish to study the time that elapses until the queue becomes empty. This is the duration of a busy period started at x , and corresponds to a first-passage time for a Wiener process which we denote by t_x . Moreover one might also be interested in the area beneath the sample path until it reaches the boundary of Q , since this area gives the total time waited by all customers involved in the busy period. We let this area be $A(x)$ and seek to compute $t_x = E(T_x)$ and $a(x) = E(A(x))$. Straightforward backward equation or martingale arguments give

$$t_x = -x/m$$

(2.10)

$$a(x) = -\frac{x^2}{2m} + \frac{\sigma^2}{2m^2} x$$

where m is negative, and m and σ^2 are given earlier. The distribution of the first passage time can also be easily calculated; however, its Laplace transform is most convenient to find and is given by

$$E(e^{-s\tau_x}) = \exp \left[\left(\frac{x}{\sigma}\right) - \left(\frac{m}{\sigma}\right) - \sqrt{\left(\frac{m}{\sigma}\right)^2 + 2s} \right] \quad (2.11)$$

4. A Refined Diffusion Approximation

In this section we continue the development of a diffusion approximation for the data queue length process. The analysis is patterned after the important semigroup methods of Burman [1979]. One develops a sequence of Markov processes and studies the behavior of their generators. One shows that the generators converge and hence concludes that the associated semi-groups converge. This entails the convergence of the finite dimensional distributions. The limiting finite dimensional distributions will be those of a certain Brownian motion. The theory underlying this approach is based on theorems of Trotter and Kato, Kurtz and Burman; the reader is referred to Burman (1979) for technical details. We will illustrate the approach in the context of the special case $v = 1, c = 0$. The method applies to the case of general c and v but the details are not given here.

Let $\{V(t), t \geq 0\}$ be the voice process, a Markov process with state space $\{0,1\}$. Let $N(t)$ be the data system size at time t . We study the sequence of Markov processes $\langle \{(X_n(t), V_n(t)), t \geq 0\} \rangle_{n=1}^{\infty}$ where

$$X_n(t) = N(nt)/\sqrt{n} \quad \text{and} \quad V_n(t) = V(nt) . \quad (3.1)$$

The generator of this bivariate process is easy to compute since $V_n(t)$ subordinates $X_n(t)$. Let $f(x,k)$ be a function with domain $[0,\infty) \times \{0,1\}$ which is smooth as a function of x for each $k = 0,1$. The generator is given by

$$A_n f(x,k) = \begin{cases} \delta n [f(x + 1/\sqrt{n}, k) - f(x,k)] \\ \quad + (1-k)\eta n [f(x - 1/\sqrt{n}, k) - f(x,k)] + nQf(x,k) \\ \qquad \qquad \qquad \text{for } x \geq 1/\sqrt{n} \\ \delta n [f(1/\sqrt{n}, k) - f(0,k)] + nQf(0,k) \\ \qquad \qquad \qquad \text{for } x = 0 . \end{cases} \quad (3.2)$$

where Q is the generator of the M/M/1/1 voice process and is given by (2.1).

Next expand in a Taylor series and collect terms to rewrite (3.2) as

$$A_n f(x,k) = \begin{cases} nQf(x,k) + \sqrt{n} f_x(x,k) (\delta - (1-k)\eta) \\ \quad + \frac{1}{2} f_{xx}(x,k) (\delta + (1-k)\eta) + o(1) \\ \qquad \qquad \qquad \text{for } x \geq 1/\sqrt{n} \\ nQf(0,k) + \sqrt{n} f_x(0,k)\delta + \frac{1}{2} f_{xx}(0,k)\delta + o(1) \\ \qquad \qquad \qquad \text{for } x = 0 \end{cases} \quad (3.3)$$

where $f_x(x,k) = \frac{\partial}{\partial x} f(x,k)$ and $f_{xx}(x,k) = \frac{\partial^2}{\partial x^2} f(x,k)$.

We wish to let $n \rightarrow \infty$ and focus attention on the data queue process alone. To derive a diffusion limit for the data

queue process alone, we introduce a sequence of functions $\langle f_n(x,k) \rangle_{n=1}^{\infty}$ for which $f_n(x,k) \rightarrow f(x)$, where f is a smooth function satisfying $f'(0) = 0$. We wish to study the limiting behavior of $A_n f_n(x,k)$. Let

$$f_n(x,k) = f(x) + \frac{1}{\sqrt{n}} g(x,k) + \frac{1}{n} h(x,k)$$

where g and h are smooth. Clearly $f_n(x,k) \rightarrow f(x)$. Substitution in (3.3) gives

$$A_n f_n(x,k) = \begin{cases} nQf(x) + \sqrt{n} [Qg(x,k) + f'(x) (\delta - (1-k)\eta)] \\ \quad + [Qh(x,k) + g_x(x,k) (\delta - (1-k)\eta) + \frac{1}{2} f''(x) (\delta + (1-k)\eta)] \\ \hspace{15em} x \geq 1/\sqrt{n} \\ \\ nQf(0) + \sqrt{n} [Qg(0,k) + f'(0)\delta] \\ \quad + [Qh(0,k) + g_x(0,k)\delta + \frac{1}{2} f''(0) (\delta)] \hspace{2em} x = 0 . \end{cases} \quad (3.4)$$

We first note that $Qf(x) = 0$ since Q operates on the voice or k component only, not on the data. This eliminates the first terms $nQf(x)$ and $nQf(0)$. We next examine the $Qg(x,k) + f'(x) (\delta - (1-k)\eta)$ term. This can be rewritten as

$$\begin{aligned}
& Qg(x,k) + f'(x) [\delta - (1-k)\eta - ((\delta-\eta) + \delta\rho_V)/(1 + \rho_V) \\
& \quad + f'(x) ((\delta-\eta) + \delta\rho_V)/(1 + \rho_V) \\
& = Qg(x,k) + f'(x) \left[-\frac{\rho_V}{1 + \rho_V} + k \right] = f'(x) \eta(1-\rho) .
\end{aligned}$$

We now select the function $g(x,k)$ so that

$$Qg(x,k) = -f'(x) \left[-\frac{\rho_V}{1 + \rho_V} + k \right] \eta .$$

The function $g(x,k)$ must satisfy the equations

$$\begin{aligned}
\lambda(g(x,1) - g(x,0)) &= f'(x) \frac{\eta\rho_V}{1 + \rho_V} \\
-\mu(g(x,1) - g(x,0)) &= -f'(x) \frac{\eta}{1 + \rho_V}
\end{aligned} \tag{3.5}$$

The above equations are consistent and redundant, thus any $g(x,k)$ for which

$$g(x,1) - g(x,0) = \frac{f'(x)\eta}{\mu(1 + \rho_V)}$$

will suffice. We select

$$\begin{aligned}
g(x,0) &= \frac{1}{2} f'(x) \\
g(x,1) &= \left(\frac{1}{2} + \frac{\eta}{\mu(1 + \rho_V)} \right) f'(x)
\end{aligned} \tag{3.6}$$

or

$$g(x,k) + \left(\frac{1}{2} + \frac{\eta k}{\mu(1 + \rho_V)} \right) f'(x) .$$

This choice of g gives

$$A_n f_n(x, k) = \begin{cases} -\sqrt{n} \eta(1-\rho) f'(x) + [Qh(x, k) + \left(\frac{1}{2} + \frac{\eta k}{\mu(1+\rho_V)}\right) f''(x) (\delta - (1-k)\eta) \\ \quad + \frac{1}{2} f''(x) (\delta + (1-k)\eta)], & x \geq 1/\sqrt{n} \\ \sqrt{n} \left(f'(0)\delta + f'(0) \frac{\eta(\rho_V - k(1+\rho_V))}{\mu(1+\rho_V)} \right) \\ \quad + \left[Qh(0, k) + \left(\frac{1}{2} + \frac{\eta k}{\mu(1+\rho_V)}\right) f''(0)\delta + \frac{1}{2} f''(0)\delta \right] & x = 0 \end{cases} \quad (3.7)$$

Equation (3.7) can be rewritten recalling $f'(0) = 0$ as

$$A_n f_n(x, k) = \begin{cases} -\sqrt{n} \eta(1-\rho) f'(x) + \left[Qh(x, k) + \eta f''(x) \left\{ \frac{1}{2} + \frac{\eta k}{\mu(1+\rho_V)} (\rho_d - (1-k)) \right. \right. \\ \quad \left. \left. + \frac{1}{2} (\rho_d + (1-k)) \right\} \right] + o(1), & x \geq 1/\sqrt{n} \\ Qh(0, k) + \eta f''(0) \left[\rho_d \frac{1}{2} + \frac{\eta k}{\mu(1+\rho_V)} + \frac{1}{2} \rho_d \right], & x = 0 \end{cases} \quad (3.8)$$

or

$$A_n f_n(x, k) = \begin{cases} -\sqrt{n} \eta(1-\rho) f'(x) + \left[Qh(x, k) + f''(x) \eta \cdot \rho_d \left(1 + \frac{\eta k}{\mu(1+\rho_V)} \right) \right] + o(1) \\ \quad x \geq 1/\sqrt{n} \\ Qh(0, k) + f''(0) \eta \rho_d \left(1 + \frac{\eta k}{\mu(1+\rho_V)} \right) + o(1), & x = 0 \end{cases} \quad (3.9)$$

We rewrite

$$Qh(x,k) + f''(x)\eta\rho_d \left[1 + \left(\frac{\eta k}{\mu(1+\rho_v)} \right) \right]$$

as

$$\phi h(x,k) + f'(x) \frac{\eta^2 \rho_d}{\mu(1+\rho_v)} \left(k - \frac{\rho_v}{1+\rho_v} \right) + f''(x)\eta\rho_d \left(\frac{\eta\rho_v}{\mu(1+\rho_v)^2} + 1 \right)$$

and choose $h(x,k)$ so that

$$Qh(x,k) = -f''(x) \frac{\eta^2 \rho_d}{\mu(1+\rho_v)} \left(k - \frac{\rho_v}{1+\rho_v} \right)$$

The function $h(x,k)$ must satisfy

$$\lambda(h(x,1) - h(x,0)) = f''(x) \frac{\eta^2 \rho_d \rho_v}{\mu(1+\rho_v)^2} \tag{3.10}$$

$$-\mu(h(x,1) - h(x,0)) = -f''(x) \frac{\eta^2 \rho_d}{\mu(1+\rho_v)^2}$$

These equations are consistent and redundant so a manifold of solutions are possible. Any one will suffice. This choice of $h(x,k)$ allows one to rewrite (3.9) as

$$A_n f_n(x, k) = \begin{cases} -\sqrt{n} \eta(1-\rho) f'(x) + f''(x) \eta \rho_d \left(\frac{\eta \rho_v}{\mu(1+\rho_v)^2} + 1 \right) + o(1) \\ \qquad \qquad \qquad x \geq 1/\sqrt{n} \\ \\ f''(0) \eta \rho_d \left(\frac{\eta \rho_v}{\mu(1+\rho_v)^2} + 1 \right) + o(1) \end{cases} \quad (3.11)$$

We now let $n \rightarrow \infty$. In order for (3.11) to converge to a sensible limit, we must invoke the heavy traffic approximation, that $\rho \rightarrow 1$. Specifically we let $\rho = \rho_n = 1 - (\theta/\sqrt{n})$ for some $\theta > 0$. Equation (3.11) becomes in the limit

$$A_n f_n(x, k) \longrightarrow \begin{cases} -\eta\theta f'(x) + f''(x) \eta \rho_d \left(\frac{\eta \rho_v}{\mu(1+\rho_v)^2} + 1 \right), & x > 0 \\ \\ f''(0) \eta \rho_d \left(\frac{\eta \rho_v}{\mu(1+\rho_v)^2} + 1 \right), & x = 0 \\ \\ f'(0) = 0 \end{cases} \quad (3.12)$$

This is the generator of a Wiener process with reflecting barrier at 0, drift of $-\eta\theta$, and

$$\frac{\sigma^2}{2} = \eta \rho_d \left(\frac{\eta \rho_v}{\mu(1+\rho_v)^2} + 1 \right) .$$

The stationary distribution for such a process is an exponential

distribution with parameter

$$\theta / \left\{ \rho_d \left(\frac{\eta \rho_v}{\mu (1 + \rho_v)^2} + 1 \right) \right\} ,$$

thus the mean queue length is given by

$$\frac{\rho_d}{\theta} \left(\frac{\eta \rho_v}{\mu (1 + \rho_v)^2} + 1 \right) .$$

It is informative to try to apply this heavy traffic result to a case in which ρ is near but less than 1. One might replace θ by $\sqrt{n} (1-\rho)$. Since we have scaled by \sqrt{n} , the stationary distribution of $N(t)$, the unscaled queue length should be approximately exponential with parameter

$$\frac{(1-\rho)}{\rho_d \left(\frac{\eta \rho_v}{\mu (1 + \rho_v)^2} + 1 \right)}$$

The mean queue length becomes

$$\frac{\rho_d}{1-\rho} \left(\frac{\eta \rho_v}{\mu (1 + \rho_v)^2} + 1 \right) .$$

This result is in exact agreement with Fischer (1977) and is therefore exact. The fluid flow approximation treats η/μ as being large, hence $[\eta \rho_v / \mu (1 + \rho_v)^2]$ is assumed to dominate 1. This gives $(\eta \rho_d \rho_v) / [\mu (1 + \rho_v)^2 (1-\rho)]$ as the fluid flow mean queue length found in Gaver and Lehoczky (1979) and in Section 2 of this paper.

The diffusion approximation thus found represents a refinement of the fluid flow diffusion approximation. Even if η/μ is large, ρ_v may be small, so the l may be important.

Once the Wiener process infinitesimal drift and variance have been found, one can also use the dynamics of the Wiener process to model the dynamics of the data queue process. Busy period distributions, areas, and transient behavior in general can be determined.

BIBLIOGRAPHY

- Barbacci, M. R. and Oakley, J. D. (1976). "The integration of Circuit and Packet Switching Networks Toward a SENET Implementation," 15th NBS-ACM Annual Technique Symposium.
- Bhat, U. N. and Fischer, M. J. (1976). "Multichannel Queueing Systems with Heterogeneous Classes of Arrivals," Naval Research Logistics Quarterly 23
- Brady, P. T. (1967). "A Statistical Analysis of On-Off Patterns in 16 Conversations." Bell S.T.J., 73-91.
- Burman, D. Y. (1979). "An Analytic Approach to Diffusion Approximations in Queueing." Ph.D. Dissertation, New York University.
- Chang, Lih-Hsing (1977). "Analysis of Integrated Voice and Data Communication Network," Ph.D. Dissertation, Department of Electrical Engineering, Carnegie-Mellon University, November.
- Coviello, G. and Vena, P. A. (1975). "Integration of Circuit/ Packet Switching in a SENET (Slotted Envelop NETWORK) Concept," National Telecommunications Conference, New Orleans, December, pp. 42-12 to 42-17.
- Fischer, M. J. (1977a). "A Queueing Analysis of an Integrated Telecommunications System with Priorities," INFOR 15
- Fischer, M. J. (1977b). "Performance of Data Traffic in an Integrated Circuit- and Packet-Switched Multiplex Structure," DCA Technical Report.
- Fischer, M. J. and Harris, T. C. (1976). "A Model for Evaluating the Performance of an Integrated Circuit- and Packet-Switched Multiplex Structure," IEEE Trans. on Comm., Com-24 February.
- Halfin, S. (1972). "Steady-state Distribution for the Buffer Content of an M/G/1 Queue with Varying Service Rate," SIAM J. Appl. Math., 356-363.
- Halfin, S. and Segal, M. (1972). "A Priority Queueing Model for a Mixture of Two Types of Customers," SIAM J. Appl. Math., 369-379.

INITIAL DISTRIBUTION LIST

	Number of Copies
Defense Documentation Center Cameron Station Alexandria, VA 22314	2
Library Code Code 0142 Naval Postgraduate School Monterey, CA 93940	2
Library Code 55 Naval Postgraduate School Monterey, Ca. 93940	1
Dean of Research Code 012A Naval Postgraduate School Monterey, Ca. 93940	1
Attn: A. Andrus, Code 55	1
D. Gaver, Code 55	25
D. Barr, Code 55	1
P. A. Jacobs, Code 55	1
P. A. W. Lewis, Code 55	1
P. Milch, Code 55	1
R. Richards, Code 55	1
M. G. Sovereign, Code 55	1
R. J. Stampfel, Code 55	1
R. R. Read, Code 55	1
J. Wozencraft, Code 74	1
Mr. Peter Badgley ONR Headquarters, Code 102B 800 N. Quincy Street Arlington, VA 22217	1
Dr. James S. Bailey, Director Geography Programs, Department of the Navy ONR Arlington, VA 93940	1
Prof. J. Lehoczky Dept. of Statistics Carnegie Mellon University Pittsburgh, PA. 15213	5

DISTRIBUTION LIST

	No. of Copies
STATISTICS AND PROBABILITY PROGRAM OFFICE OF NAVAL RESEARCH CODE 422 ARLINGTON VA	1 22217
OFFICE OF NAVAL RESEARCH NEW YORK AREA OFFICE 715 BROADWAY - 5TH FLOOR ATTN: DR. ROBERT GRAFTON NEW YORK, NY	1 10003
DIRECTOR OFFICE OF NAVAL RESEARCH BRANCH OFF 536 SOUTH CLARK STREET ATTN: DEPUTY AND CHIEF SCIENTIST CHICAGO, IL	1 60605
LIBRARY NAVAL OCEAN SYSTEMS CENTER SAN DIEGO CA	1 92152
NAVY LIBRARY NATIONAL SPACE TECHNOLOGY LAB ATTN: NAVY LIBRARIAN BAY ST. LOUIS MS	1 39522
NAVAL ELECTRONIC SYSTEMS COMMAND NAVELEX 320 NATIONAL CENTER NO. 1 ARLINGTON VA	1 20360
DIRECTOR NAVAL RESEARCH LABORATORY ATTN: LIBRARY (DNRL) CODE 2025 WASHINGTON, D.C.	1 20375
TECHNICAL INFORMATION DIVISION NAVAL RESEARCH LABORATORY WASHINGTON, D. C.	1 20375

DISTRIBUTION LIST

No. of Copies

PROF. C. R. BAKER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL, NORTH CAROLINA 27514	1
PROF. R. E. DECHOFER DEPARTMENT OF OPERATIONS RESEARCH CORNELL UNIVERSITY ITHACA NEW YORK 14850	1
PROF. N. J. BERSHAD SCHOOL OF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE, CALIFORNIA 92664	1
P. J. BICKEL DEPARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA BERKELEY, CALIFORNIA 94720	1
PROF. F. W. BLOCK DEPARTMENT OF MATHEMATICS UNIVERSITY OF PITTSBURGH PITTSBURGH PA 15260	1
PROF. JOSEPH BLUM DEPT. OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE THE AMERICAN UNIVERSITY WASHINGTON DC 20016	1
PROF. R. A. BRADLEY DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE, FLORIDA 32306	1
PROF. R. E. BARLOW OPERATIONS RESEARCH CENTER COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BERKLEY CALIFORNIA 94720	1
MR. C. N. BENNETT NAVAL COASTAL SYSTEMS LABORATORY CODE P701 PANAMA CITY, FLORIDA 32401	1

DISTRIBUTION LIST

No. of Copies

PROF. L. N. PHAT COMPUTER SCIENCE / OPERATIONS RESEARCH CENTER SOUTHERN METHODIST UNIVERSITY DALLAS TEXAS 75275	1
PROF. W. R. ELISCHKE DEPT. OF QUANTITATIVE BUSINESS ANALYSIS UNIVERSITY OF SOUTHERN CALIFORNIA LOS ANGELES, CALIFORNIA 90007	1
DR. DERRILL J. BERDELON NAVAL UNDERWATER SYSTEMS CENTER CODE 21 NEWPORT RI 02840	1
J. E. BOYER JR DEPT. OF STATISTICS SOUTHERN METHODIST UNIVERSITY DALLAS TX 75275	1
DR. J. CHANDRA U. S. ARMY RESEARCH P. O. BOX 12211 RESEARCH TRIANGLE PARK NORTH CAROLINA 27706	1
PROF. F. CHERNOFF DEPT. OF MATHEMATICS MASS INSTITUTE OF TECHNOLOGY CAMBRIDGE, MASSACHUSETTS 02139	1
PROF. C. GERMAN DEPARTMENT OF CIVIL ENGINEERING AND ENGINEERING MECHANICS COLUMBIA UNIVERSITY NEW YORK NEW YORK 10027	1
PROF. R. L. DISNEY VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEPT. OF INDUSTRIAL ENGINEERING AND OPERATIONS RESEARCH BLACKSBURG, VA 24061	1

DISTRIBUTION LIST

MR. GENE F. GLEISSNER 1
 APPLIED MATHEMATICS LABORATORY
 CAROL TAYLOR NAVAL SHIP RESEARCH
 AND DEVELOPMENT CENTER
 BETHESDA
 MD 20084

PROF. S. S. GUPTA 1
 DEPARTMENT OF STATISTICS
 PURDUE UNIVERSITY
 LAFAYETTE
 INDIANA 47907

PROF. C. L. HANSON 1
 DEPT OF MATH. SCIENCES
 STATE UNIVERSITY OF NEW YORK,
 BINGHAMTON
 BINGHAMTON
 NY 13901

Prof. M. J. Hinich 1
 Dept. of Economics
 Virginia Polytechnic Institute
 and State University
 Blacksburg, VA 24061

Dr. D. Depriest, 1
 ONR, Code 102B
 800 N. Quincy Street
 Arlington, VA 22217

Prof. G. E. Whitehouse 1
 Dept. of Industrial Engineering
 Lehigh University
 Bethlehem, PA 18015

Prof. M. Zia-Hassan 1
 Dept. of Ind. & Sys. Eng.
 Illinois Institute of Technology
 Chicago, IL 60616

Prof. S. Zacks 1
 Statistics Dept.
 Virginia Polytechnic Inst.
 Blacksburg, VA 24061

Head, Math. Sci Section 1
 National Science Foundation
 1800 G Street, N.W.
 Washington, D.C. 20550

No. of Copies

Dr. H. Sittrop
Physics Lab., TNO
P.O. Box 96964
2509 JG, The Hague
The Netherlands 1

DR. R. ELASHOFF
BIOMATHEMATICS
UNIV. OF CALIF.
LOS ANGELES
CALIFORNIA 1

90024

PROF. GEORGE S. FISHMAN
UNIV. OF NORTH CAROLINA
CUR. IN OR AND SYS. ANALYSIS
PHILLIPS ANNEX
CHAPEL HILL, NORTH CAROLINA 1

20742

DR. R. GRANACESIKAN
BELL TELEPHONE LAB
MOLDEL, N. J. 1

07733

DR. A. J. COLEMAN
CHIEF, CR
DIV. 205.02, ADMIN. A428
U.S. DEPT. OF COMMERCE
WASHINGTON, D.C. 1

20234

DR. H. HIGGINS
53 BONN 1, POSTFACH 589
NASSESTRASSE 2 1

WEST GERMANY

DR. P. T. HOLMES
DEPT. OF MATH.
CLEMSON UNIV.
CLEMSON
SOUTH CAROLINA 1

29631

Dr. J. A. Hocke
Bell Telephone Labs
Whippany, New Jersey 07733 1

Dr. Robert Hooke
Box 1982
Pinehurst, No. Carolina 28374 1

DR. D. L. IGLEHART
DEPT. OF C.F.
STANFORD UNIV.
STANFORD
CALIFORNIA

94305

1

Dr. D. Trizna, Mail Code 5323
Naval Research Lab
Washington, D.C. 20375

1

Dr. E. J. Wegman,
ONR, Cdoe 436
Arlington, VA 22217

1

DR. H. KOBAYASHI
IBM
YORKTOWN HEIGHTS
NEW YORK

10598

1

DR. JOHN LEHOCZKY
STATISTICS DEPARTMENT
CARNEGIE-MELLON UNIVERSITY
PITTSBURGH
PENNSYLVANIA

15213

1

DR. A. LEMOINE
1020 GUINCA ST.
PALO ALTO,
CALIFORNIA

94301

1

DR. J. MACCUSEN
UNIV. OF CALIF.
LOS ANGELES
CALIFORNIA

90024

1

FRFC. K. T. MARSHALL
DEPT. OF C.F.
NAVAL POSTGRADUATE SCHOOL
MONTEREY
CALIFORNIA

93940

1

DR. M. MAZUMCAR
MATH. DEPT.
ESTINGHOUSE RES. LABS
CHURCHILL BCF
PITTSBURGH
PENNSYLVANIA

15235

1

DISTRIBUTION LIST

No. of Copies

<p>PROF. W. M. FIRSCH INSTITUTE OF MATHEMATICAL SCIENCES NEW YORK UNIVERSITY NEW YORK NEW YORK 10453</p>	<p>1</p>
<p>PROF. J. B. KADANE DEPARTMENT OF STATISTICS CARNEGIE-MELLON PITTSBURGH, PENNSYLVANIA 15213</p>	<p>1</p>
<p>DR. RICHARD LAU DIRECTOR OFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET PASADENA CA</p>	<p>1</p>
<p>DR. A. R. LAUFER DIRECTOR OFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET PASADENA CA</p>	<p>1</p>
<p>PROF. M. LEADBETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NORTH CAROLINA 27514</p>	<p>1</p>
<p>DR. J. S. LEE J. S. LEE ASSOCIATES, INC. 2001 JEFFERSON DAVIS HIGHWAY SUITE 802 ARLINGTON VA</p>	<p>1</p>
<p>PROF. L. D. LEE DEPARTMENT OF STATISTICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSBURG VA</p>	<p>1</p>
<p>PROF. R. S. LEVENWORTH DEPT. OF INDUSTRIAL AND SYSTEMS ENGINEERING UNIVERSITY OF FLORIDA GAINESVILLE, FLORIDA 32611</p>	<p>1</p>

DISTRIBUTION LIST

No. of copies

<p>FRCF G. LIEPERMAN STANFORD UNIVERSITY DEPARTMENT OF OPERATIONS RESEARCH STANFORD CALIFORNIA 94305</p>	<p>1</p>
<p>DR. JAMES R. MAAR NATIONAL SECURITY AGENCY FORT MEADE, MARYLAND 20755</p>	<p>1</p>
<p>FPCF. R. W. MAESEN DEPARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA MO</p>	<p>1</p>
<p>65201</p>	
<p>DR. N. R. MANN SCIENCE CENTER ROCKWELL INTERNATIONAL CORPORATION P.O. BOX 1085 THOUSAND OAKS, CALIFORNIA 91320</p>	<p>1</p>
<p>DR. W. H. MARLOW PROGRAM IN LOGISTICS THE GEORGE WASHINGTON UNIVERSITY 707 22ND STREET, N. W. WASHINGTON, D. C. 20037</p>	<p>1</p>
<p>PROF. E. MASRY DEPT. APPLIED PHYSICS AND INFORMATION SERVICE UNIVERSITY OF CALIFORNIA LA JOLLA CALIFORNIA</p>	<p>1</p>
<p>92093</p>	
<p>DR. BRUCE J. MCCONALD SCIENTIFIC DIRECTOR SCIENTIFIC LIAISON GROUP OFFICE OF NAVAL RESEARCH AMERICAN EMBASSY - TOKYO AFC SAN FRANCISCO</p>	<p>1</p>
<p>96503</p>	

Dr. Leon F. McGinnis School of Ind. And Sys. Eng. Georgia Inst. of Tech. Atlanta, GA 30332	1
DR. D. R. MCNEIL DEPT. OF STATISTICS PRINCETON UNIV. PRINCETON NEW JERSEY	1 08540
DR. F. MOSTELLER STAT. DEPT. HARVARD UNIV. CAMBRIDGE MASSACHUSETTS	1 02139
DR. H. REISER IBM THOMAS J. WATSON RES. CTR. YORKTOWN HEIGHTS NEW YORK	1 10598
DR. J. RICREAN DEPT. OF MATHEMATICS ROCKEFELLER UNIV. NEW YORK NEW YORK	1 10021
DR. LINUS SCHRAGE UNIV. OF CHICAGO GRAD. SCHOOL OF BUS. 5826 GREENWOOD AVE. CHICAGO, ILLINOIS	1 60637
Dr. Paul Schweitzer University of Rochester Rochester, N.Y. 14627	1
Dr. V. Srinivasan Graduate School of Business Stanford University Stanford, CA. 94305	1
Dr. Roy Welsch M.I.T. Sloan School Cambridge, MA 02139	1

DISTRIBUTION LIST

No. of Copies

DR. JANET M. MYHRE
THE INSTITUTE OF DECISION SCIENCE
FOR BUSINESS AND PUBLIC POLICY
CLAREMONT MEN'S COLLEGE
CLAREMONT
CA 91711

1

MR. F. NISSELSOHN
BUREAU OF THE CENSUS
ROOM 2029
FEDERAL BUILDING 3
WASHINGTON,
D. C. 20333

1

MISS B. S. CRLEANS
NAVAL SEA SYSTEMS COMMAND
(SEA O3F)
RM 105C3
ARLINGTON VIRGINIA 20360

1

PROF. C. E. OWEN
DEPARTMENT OF STATISTICS
SOUTHERN METHODIST UNIVERSITY
DALLAS
TEXAS
75222

1

Prof. E. Parzen
Statistical Science Division
Texas A & M University
College Station TX 77843

1

DR. A. PETRASOVITS
RCCM 207E, FCCC AND ERIG BLDG.
TUNNEY'S PASTURE
CITOWA, ONTARIO K1A-CL2,
CANADA

1

PROF. S. L. PFCENIX
SIBLEY SCHOOL OF MECHANICAL AND
AEROSPACE ENGINEERING
CORNELL UNIVERSITY
ITHACA
NY 14850

1

DR. A. L. POWELL
DIRECTOR
OFFICE OF NAVAL RESEARCH BRANCH OFF
495 SUMMER STREET
BOSTON
MA 02210

1

MR. F. R. FRICFI
CODE 224 OPERATIONAL TEST AND ONRS
EVALUATION FORCE (OPTVEFOR)
NORFOLK,
VIRGINIA
20360

1

DISTRIBUTION LIST

No. of Copies

PROF. M. L. PURI DEPT. OF MATHEMATICS P.O. BOX F INDIANA UNIVERSITY FOUNDATION BLOOMINGTON IN 47401	1
PROF. H. ROBBINS DEPARTMENT OF MATHEMATICS COLUPEIA UNIVERSITY NEW YORK, NEW YORK 10027	1
PROF. M. ROSENBLATT DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA 92093	1
PROF. S. M. ROSS COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BERKELEY CA 94720	1
PROF. I. RUBIN SCHOOL OF ENGINEERING AND APPLIED SCIENCE UNIVERSITY OF CALIFORNIA LOS ANGELES, CALIFORNIA 90024	1
PROF. I. R. SAVAGE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN, CONNECTICUT 06520	1
PROF. L. L. SCHARF JR DEPARTMENT OF ELECTRICAL ENGINEERING COLORADO STATE UNIVERSITY FT. COLLINS, COLORADO 80521	1
PROF. R. SERFLING DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE FLORIDA 32306	1
PROF. W. R. SCHUCANY DEPARTMENT OF STATISTICS SOUTHERN METHODIST UNIVERSITY DALLAS, TEXAS 75222	1

DISTRIBUTION LIST

No. of Copies

PROF. C. C. SIEGMUND DEPT. OF STATISTICS STANFORD UNIVERSITY STANFORD CA	1
54305	
PROF. M. L. SLOMAN DEPT. OF ELECTRICAL ENGINEERING POLYTECHNIC INSTITUTE OF NEW YORK BROOKLYN, NEW YORK 11201	1
DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANDANT OF THE MARINE CORPS WASHINGTON, D. C. 20380	1
DR. C. E. SMITH DESMATICS INC. P.O. BOX 618 STATE COLLEGE PENNSYLVANIA 16801	1
PROF. W. L. SMITH DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NORTH CAROLINA 27514	1
Dr. H. J. Solomon ONR 223/231 Old Marylebone Rd London NW1 5TH, ENGLAND	1
MR. GLENN F. STAFFLY NATIONAL SECURITY AGENCY FORT MEADE, MARYLAND 20755	1
Mr. J. Gallagher Naval Underwater Systems Center New London, CT	1
Dr. E. C. Monahan Dept. of Oceanography University College Galway, Ireland	1

DISTRIBUTION LIST

No. of Copies

DR. R. M. STARK STATISTICS AND COMPUTER SCI. UNIV. OF DELAWARE NEWARK DELAWARE	19711	1
PROF. RICHARD VANSLYKE RES. ANALYSIS CORP. BEECHWOOD CLD TAPPEN ROAD GLEN COVE, NEW YORK	11542	1
PROF. JOHN W. TUKEY FINE HALL PRINCETON UNIV. PRINCETON NEW JERSEY	08540	1
DR. THOMAS C. VARLEY OFFICE OF NAVAL RESEARCH CODE 424 ARLINGTON VA	22217	1
PROF. G. WATSON FINE HALL PRINCETON UNIV. PRINCETON NEW JERSEY	08540	1
MR. DAVID A. SWICK ADVANCED PROJECTS GROUP CODE 8103 NAVAL RESEARCH LAB. WASHINGTON DC	20375	1
MR. WENDELL G. SYKES ARTHUR C. LITTLE, INC. ACORN PARK CAMBRIDGE MA	02140	1
PROF. J. R. THOMPSON DEPARTMENT OF MATHEMATICAL SCIENCE RICE UNIVERSITY HOUSTON, TEXAS 77001		1
PROF. W. A. THOMPSON DEPARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA, MISSOURI 65201		1

DISTRIBUTION LIST

	No. of Copies
PROF. F. A. TILLMAN DEPT. OF INDUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN KS	1
66506	
PROF. J. W. TUKEY DEPARTMENT OF STATISTICS PRINCETON UNIVERSITY PRINCETON, N. J. 08540	1
PROF. A. F. VEINOTT DEPARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD CALIFORNIA 94305	1
DANIEL H. WAGNER STATION SQUARE ONE FACIL, PENNSYLVANIA 19301	1
PROF. GRACE WAHBA DEPT. OF STATISTICS UNIVERSITY OF WISCONSIN MADISON WI	1
53706	
PROF. K. T. WALLENIS DEPARTMENT OF MATHEMATICAL SCIENCES, CLEMSON UNIVERSITY CLEMSON, SOUTH CAROLINA 29631	1
PROF. BERNARD WIDROW STANFORD ELECTRONICS LAB STANFORD UNIVERSITY STANFORD CA	1
94305	

DISTRIBUTION LIST

No. of Copies

OFFICE OF NAVAL RESEARCH SAN FRANCISCO AREA OFFICE 760 MARKET STREET SAN FRANCISCO CALIFORNIA 94102	1
TECHNICAL LIBRARY NAVAL ORDNANCE STATION INDIAN HEAD MARYLAND 20640	1
NAVAL SHIP ENGINEERING CENTER PHILADELPHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19112	1
BUREAU OF NAVAL PERSONNEL DEPARTMENT OF THE NAVY TECHNICAL LIBRARY WASHINGTON D. C. 20370	1
PROF. M. AECHE-HAMEZD DEPARTMENT OF MATHEMATICS UNIVERSITY OF NORTH CAROLINA CHARLOTTE NC 28223	1
PROF. T. W. ANDERSON DEPARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORD , CALIFORNIA 94305	1
PROF. F. J. ANSCOMBE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN CONNECTICUT 06520	1
PROF. L. A. ARCIA INSTITUTE OF INDUSTRIAL ADMINISTRATION UNION COLLEGE SCHENECTADY , NEW YORK 12308	1

U190212

DUDLEY KNOX LIBRARY - RESEARCH REPORTS



5 6853 01069331 0

019021